

# Large-step semantics

CS 152 (Spring 2021)

Harvard University

Thursday, February 4, 2021

# Small-step semantics (review)

$\longrightarrow \subseteq \mathbf{Config} \times \mathbf{Config}$

$\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$

# Large-step semantics

$$\Downarrow \subseteq \mathbf{Config} \times \mathbf{FinalConfig}$$

where

$$\mathbf{Config} = \mathbf{Exp} \times \mathbf{Store}$$

$$\text{and } \mathbf{Final Config} = \mathbf{Int} \times \mathbf{Store} \subseteq \mathbf{Config}.$$

We write  $\langle e, \sigma \rangle \Downarrow \langle n, \sigma' \rangle$  to mean that  
 $(\langle e, \sigma \rangle, \langle n, \sigma' \rangle) \in \Downarrow$

# Semantic Rules (1)

$$\text{INT}_{\text{LRG}} \frac{}{\langle n, \sigma \rangle \Downarrow \langle n, \sigma \rangle}$$

$$\text{VAR}_{\text{LRG}} \frac{}{\langle x, \sigma \rangle \Downarrow \langle n, \sigma \rangle} \text{ where } \sigma(x) = n$$

$$\text{ADD}_{\text{LRG}} \frac{\begin{array}{l} \langle e_1, \sigma \rangle \Downarrow \langle n_1, \sigma'' \rangle \\ \langle e_2, \sigma'' \rangle \Downarrow \langle n_2, \sigma' \rangle \\ \text{where } n \text{ is the sum of } n_1 \text{ and } n_2 \end{array}}{\langle e_1 + e_2, \sigma \rangle \Downarrow \langle n, \sigma' \rangle}$$

## Semantic Rules (2)


$$\text{MUL}_{\text{LRG}} \frac{\begin{array}{l} \langle e_1, \sigma \rangle \Downarrow \langle n_1, \sigma'' \rangle \\ \langle e_2, \sigma'' \rangle \Downarrow \langle n_2, \sigma' \rangle \end{array} \text{ where } n \text{ is the product of } n_1 \text{ and } n_2}{\langle e_1 \times e_2, \sigma \rangle \Downarrow \langle n, \sigma' \rangle}$$

$$\text{ASG}_{\text{LRG}} \frac{\begin{array}{l} \langle e_1, \sigma \rangle \Downarrow \langle n_1, \sigma'' \rangle \\ \langle e_2, \sigma''[x \mapsto n_1] \rangle \Downarrow \langle n_2, \sigma' \rangle \end{array}}{\langle x := e_1; e_2, \sigma \rangle \Downarrow \langle n_2, \sigma' \rangle}$$

# Example

$\langle \text{foo} := 3; \text{foo} \times \text{bar}, \sigma \rangle \Downarrow \langle 21, \sigma' \rangle$  for a store  $\sigma$  such that  $\sigma(\text{bar}) = 7$ , and  $\sigma' = \sigma[\text{foo} \mapsto 3]$ .

# Example Proof Tree



# Observation

a depth-first traversal of the large-step proof tree yields the sequence of one-step transitions in small-step evaluation.



# Equivalence of semantics

For all expressions  $e$ , stores  $\sigma$  and  $\sigma'$ , and integers  $n$ , we have:

$$\langle e, \sigma \rangle \Downarrow \langle n, \sigma' \rangle \iff \langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle .$$



$$P(e) = \forall \sigma, \sigma' \in \mathbf{Store.}, \forall n \in \mathbf{Int.}$$

$$\langle e, \sigma \rangle \Downarrow \langle n, \sigma' \rangle$$



$$\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$$

Proof by structural induction on expression  $e$ .

$\implies$  Case  $e \equiv x$

$\implies$  Case  $e \equiv x$

- ▶ By hypothesis,  $\langle x, \sigma \rangle \Downarrow \langle n, \sigma' \rangle$ .
- ▶ By inversion,  $n = \sigma(x)$  and  $\sigma = \sigma'$ .
- ▶ By VAR,  $\langle x, \sigma \rangle \longrightarrow \langle n, \sigma \rangle$ .
- ▶ We can conclude  $\langle x, \sigma \rangle \longrightarrow^* \langle n, \sigma \rangle$ .

$\implies$  Case  $e \equiv n$

$\implies$  Case  $e \equiv n$

- ▶ By hypothesis and inversion, we know that

$$\langle n, \sigma \rangle \Downarrow \langle n, \sigma \rangle$$

- ▶ By trivial reflexivity

$$\langle n, \sigma \rangle \longrightarrow^* \langle n, \sigma \rangle$$

$\implies$  Case  $e \equiv e_1 + e_2$

$\implies$  Case  $e \equiv e_1 + e_2$

$$P(e_1) = \forall n, \sigma, \sigma' : \quad \langle e_1, \sigma \rangle \Downarrow \langle n, \sigma' \rangle \implies \\ \langle e_1, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$$

$$P(e_2) = \forall n, \sigma, \sigma' : \quad \langle e_2, \sigma \rangle \Downarrow \langle n, \sigma' \rangle \implies \\ \langle e_2, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$$

$$P(e) = \forall n, \sigma, \sigma' : \quad \langle e_1 + e_2, \sigma \rangle \Downarrow \langle n, \sigma' \rangle \implies \\ \langle e_1 + e_2, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$$



# Lemma 1

If  $\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$  then for all  $n_1, e_2$  the following hold.

▶  $\langle e + e_2, \sigma \rangle \longrightarrow^* \langle n + e_2, \sigma' \rangle$

▶  $\langle e \times e_2, \sigma \rangle \longrightarrow^* \langle n \times e_2, \sigma' \rangle$

▶  $\langle n_1 + e, \sigma \rangle \longrightarrow^* \langle n_1 + n, \sigma' \rangle$

▶  $\langle n_1 \times e, \sigma \rangle \longrightarrow^* \langle n_1 \times n, \sigma' \rangle$

$\implies$  Case  $e \equiv e_1 \times e_2$

$\implies$  Case  $e \equiv x := e_1; e_2$

Need lemma similar to lemma1:

$\forall e \in \mathbf{Exp}. \forall \sigma, \sigma' \in \mathbf{Store}, \forall n \in \mathbf{Int}.$   
if  $\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$  then  
 $\forall x \in \mathbf{Var}. \forall e_2 \in \mathbf{Exp}.$   
 $\langle x := e; e_2, \sigma \rangle \longrightarrow^* \langle e_2, \sigma'[x \mapsto n] \rangle .$



Proof by mathematical induction on the number of steps  $\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$ .

⇐ Base case

## ⇐ Base case

If  $\langle e, \sigma \rangle \xrightarrow{*} \langle n, \sigma' \rangle$  in zero steps, then we must have  $e \equiv n$  and  $\sigma' = \sigma$ . By  $\text{INT}_{\text{LRG}}$ ,  
 $\langle n, \sigma \rangle \Downarrow \langle n, \sigma \rangle$ .

⇐ Inductive case

## ⇐ Inductive case

- ▶ Assume  $\langle e, \sigma \rangle \rightarrow \langle e'', \sigma'' \rangle \rightarrow^* \langle n, \sigma' \rangle$ .
- ▶ Inductive Hypothesis:  $\langle e'', \sigma'' \rangle \Downarrow \langle n, \sigma' \rangle$ .



## Lemma 2

For all  $e, e', \sigma$ , and  $n$ , if  $\langle e, \sigma \rangle \longrightarrow \langle e', \sigma'' \rangle$  and  $\langle e', \sigma'' \rangle \Downarrow \langle n, \sigma' \rangle$ , then  $\langle e, \sigma \rangle \Downarrow \langle n, \sigma' \rangle$ .

# Lemma 1

If  $\langle e, \sigma \rangle \longrightarrow^* \langle n, \sigma' \rangle$  then for all  $n_1, e_2$  the following hold.

▶  $\langle e + e_2, \sigma \rangle \longrightarrow^* \langle n + e_2, \sigma' \rangle$

▶  $\langle e \times e_2, \sigma \rangle \longrightarrow^* \langle n \times e_2, \sigma' \rangle$

▶  $\langle n_1 + e, \sigma \rangle \longrightarrow^* \langle n_1 + n, \sigma' \rangle$

▶  $\langle n_1 \times e, \sigma \rangle \longrightarrow^* \langle n_1 \times n, \sigma' \rangle$

# Proof of Lemma1

By mathematical induction on the number of evaluation steps in  $\longrightarrow^*$ .

## Lemma 2

For all  $e, e', \sigma$ , and  $n$ , if  $\langle e, \sigma \rangle \longrightarrow \langle e', \sigma'' \rangle$  and  $\langle e', \sigma'' \rangle \Downarrow \langle n, \sigma' \rangle$ , then  $\langle e, \sigma \rangle \Downarrow \langle n, \sigma' \rangle$ .

# Proof of Lemma 2

(part of homework)

# Break: Easier in Small-step or Large-step?

- ▶ Proving totality
- ▶ Dealing with non-termination
- ▶ Expressing interleaving / concurrency
- ▶ Your ideas?

# Scratchpad

# Stuck vs Non-Terminating

$$e ::= \dots$$

- | stop
- | loop

$$\text{LOOP} \frac{}{\langle \text{loop}, \sigma \rangle \longrightarrow \langle \text{loop}, \sigma \rangle}$$
$$\text{LOOP}_{\text{LRG}} \frac{\langle \text{loop}, \sigma \rangle \Downarrow \langle n, \sigma' \rangle}{\langle \text{loop}, \sigma \rangle \Downarrow \langle n, \sigma' \rangle}$$



# Rules Add (Original)

$$\text{LADD} \frac{\langle e_1, \sigma \rangle \longrightarrow \langle e'_1, \sigma' \rangle}{\langle e_1 + e_2, \sigma \rangle \longrightarrow \langle e'_1 + e_2, \sigma' \rangle}$$

$$\text{RADD} \frac{\langle e_2, \sigma \rangle \longrightarrow \langle e'_2, \sigma' \rangle}{\langle n + e_2, \sigma \rangle \longrightarrow \langle n + e'_2, \sigma' \rangle}$$

$$\text{ADD} \frac{\text{where } p \text{ is the sum of } n \text{ and } m}{\langle n + m, \sigma \rangle \longrightarrow \langle p, \sigma \rangle}$$

# Rules Add (Interleaving)

$$\text{LADD} \frac{\langle e_1, \sigma \rangle \longrightarrow \langle e'_1, \sigma' \rangle}{\langle e_1 + e_2, \sigma \rangle \longrightarrow \langle e'_1 + e_2, \sigma' \rangle}$$

$$\text{RADD} \frac{\langle e_2, \sigma \rangle \longrightarrow \langle e'_2, \sigma' \rangle}{\langle e_1 + e_2, \sigma \rangle \longrightarrow \langle e_1 + e'_2, \sigma' \rangle}$$

$$\text{ADD} \frac{\text{where } p \text{ is the sum of } n \text{ and } m}{\langle n + m, \sigma \rangle \longrightarrow \langle p, \sigma \rangle}$$